

On lower bounds of the solutions of some simple reaction-diffusion equations

Malte Henkel^{a,b}

^aLaboratoire de Physique des Matériaux,¹ Université Henri Poincaré Nancy I,
B.P. 239, F – 54506 Vandœuvre lès Nancy Cedex, France²

^bCentro de Física Teórica e Computacional, Universidade de Lisboa,
Av. Prof. Gama Pinto 2, P-1649-003 Lisboa, Portugal

Abstract

The mean-field reaction-diffusion equations of the diffusive pair-annihilation and triplett-annihilation processes are considered. A direct lower bound on the time-dependent mean particle-density is derived. The results are applied to the mean-field theory of the diffusive pair-contact process.

PACS numbers: 05.10.Gg, 05.70.ln, 02.30.Jr

¹Laboratoire associé au CNRS UMR 7556

²permanent address

1 Introduction

The interplay of reaction kinetics with particle diffusion has since a long time been a topic of intensive research. Here we shall consider the following reaction-diffusion equations

$$\partial_t a(t, \mathbf{r}) = \Delta a(t, \mathbf{r}) - \lambda a(t, \mathbf{r})^2 \quad (1.1)$$

and

$$\partial_t b(t, \mathbf{r}) = \Delta b(t, \mathbf{r}) - \mu b(t, \mathbf{r})^3 \quad (1.2)$$

where $t \in \mathbb{R}_+$ and $\mathbf{r} \in \mathbb{R}^n$ are time and space coordinates, Δ is the Laplacian and $\lambda, \mu > 0$ are constant reaction rates. It is well-known that eq. (1.1) provides a mean-field description of the pair-annihilation process $A + A \rightarrow \emptyset$ together with single-particle diffusion and similarly, (1.2) describes triplet annihilation $A + A + A \rightarrow \emptyset$ (we have rescaled the diffusion-constant to one). Still, the derivation of the long-time behaviour of solutions of such non-linear partial differential equations is not completely trivial. It is convenient to consider eq. (1.1) inside a spatial domain $\Omega \subset \mathbb{R}^n$ and to define the mean densities

$$\bar{a} = \bar{a}(t) = \frac{1}{|\Omega|} \int_{\Omega} d\mathbf{r} a(t, \mathbf{r}) \quad ; \quad \bar{b} = \bar{b}(t) = \frac{1}{|\Omega|} \int_{\Omega} d\mathbf{r} b(t, \mathbf{r}) \quad (1.3)$$

where $|\Omega|$ denotes the volume of Ω .

The most simplistic treatment of (1.1) simply suppresses the diffusion term, which leads to $\partial_t \bar{a}_s = -\lambda \bar{a}_s^2$ and the solution $\bar{a}_s(t) = a_0/(1 + a_0 \lambda t)$. Then, it may be asked to what extent this drastic simplification may be justified. This, and more generally the long-time behaviour of the space-time-dependent particle density $a(t, \mathbf{r})$, has received a lot of attention. For example, *a priori* estimates such as the strong maximum principle may be invoked to obtain bounds $v_-(t) \leq a(t, \mathbf{r}) \leq v_+(t)$ where $v_{\pm}(t)$ satisfy the simplistic equation $\partial_t v_{\pm} = -\lambda v_{\pm}^2$ with the initial conditions $v_-(0) \leq a(0, \mathbf{r}) \leq v_+(0)$ [12, p. 94]. The latter conditions might however be difficult to meet for very ‘rough’ initial data $a(0, \mathbf{r})$. For domains with a finite (and ‘small’) volume $|\Omega|$, one may define an invariant region Σ and define $\sigma := \eta - M$, where η is the principal eigenvalue of $-\Delta$ on Ω and $M = \max_{\Sigma} |\nabla \lambda a^2|$. If $\sigma > 0$, then it can be shown that $a(t, \mathbf{r})$ converges exponentially fast with a characteristic time $1/\sigma$ towards the solution $\bar{a}_s(t)$ [12, p. 223]. However, the implied exponential approach need not hold any longer in spatially infinite regions. Then methods based on a scaling *ansatz* of the form $a(t, \mathbf{r}) = t^{-\alpha/2} f(\mathbf{r} t^{-1/2})$ permit to extract the long-time asymptotics of the solution from phenomenological scaling [5] or by using rigorous renormalization-group arguments [3]. This kind of argument can also be extended to systems of reaction-diffusion equations and in particular the reactions fronts in the two-component system $kA + kB \rightarrow \emptyset$ with initial conditions such that there is a reaction front between A-rich and B-rich regions. In $n = 1$ dimension, it was shown rigorously for $k = 1$ [10] and $k \geq 4$ [2] that the particle densities and the reaction front satisfy a multiscaling behaviour and that the convergence towards the scaling solutions is controlled by algebraically (and not exponentially) small corrections in t . On the diffusive scale, when $|\mathbf{r}|/\sqrt{t} \gg 1$, the problem essentially reduces to $\partial_t a = \Delta a - a^k$ [2]. Reaction-diffusion systems of the form $\partial_t \mathbf{a} = \Delta \mathbf{a} - \nu \mathbf{a} - \mathbf{f}(\mathbf{a}) + \mathbf{g}$ with $\mathbf{a}|_{t=0} = \mathbf{a}_{(0)}$, $\mathbf{a}|_{\partial\Omega} = \mathbf{0}$, $\nu > 0$ and suitable assumptions on $\mathbf{f}$ are reviewed in [13]. The long-time behaviour of the unique solution (roughly, on the Sobolev space $W^{2,q}(\Omega)$ where $\mathbf{g} \in L^q(\Omega)$ with $q > \max(2, n/2)$) is described in terms of an attractor whose complexity can be analysed through its Kolmogorov entropy in great detail.

Here we wish to present a simple direct estimate on the mean densities $\bar{a}(t)$ and $\bar{b}(t)$. We have (see section 2 for notations)

Theorem: (i) Let $a \in C^1(\mathbb{R}_+; W^{2,2}(\Omega))$ be an (almost) everywhere non-negative solution of (1.1). Let in addition be $\nabla a = 0$ on the boundary $\partial\Omega$. Then there is a positive and $|\Omega|$ -independent constant λ' such that the mean density $\bar{a}$ satisfies

$$-\lambda' \bar{a}(t)^2 \leq \partial_t \bar{a}(t) \leq -\lambda \bar{a}(t)^2 \quad (1.4)$$

(ii) If $b \in C^1(\mathbb{R}_+; W^{2,3}(\Omega))$ is an (almost) everywhere non-negative solution of (1.2) such that $\nabla b|_{\partial\Omega} = 0$, there is a positive and $|\Omega|$ -independent constant μ' such that

$$-\mu' \bar{b}(t)^3 \leq \partial_t \bar{b}(t) \leq -\mu \bar{b}(t)^3 \quad (1.5)$$

For $n \leq 3$, the condition $b \in C^1(\mathbb{R}_+; W^{2,2}(\Omega) \cap L^3(\Omega))$ is sufficient.

The Sobolev space $W^{2,p}(\Omega)$ may be too restrictive for unbounded domains. In section 2 we define a generalized space $\widetilde{W}^{2,p}(\Omega)$ for which the theorem still holds and which includes spatially homogeneous initial states.

The upper bound in (1.4) has been known since a long time [1]. In particular, (1.4,1.5) imply for t sufficiently large (i.e. $a_0 \lambda' t > 1$ and $2b_0^2 \mu' t > 1$, respectively, where $a_0 > 0$ and $b_0 > 0$ are the initial mean densities) the bounds

$$\frac{1}{2\lambda'} \leq t \bar{a}(t) \leq \frac{1}{\lambda} \quad ; \quad \frac{1}{2\sqrt{\mu'}} \leq t^{1/2} \bar{b}(t) \leq \frac{1}{\sqrt{2\mu}} \quad (1.6)$$

It is admissible to take the infinite-volume limit $|\Omega| \rightarrow \infty$. Eq. (1.6) is in full agreement with the results established earlier by different means [2, 3, 5, 7, 12]. For similar upper bounds on $\|a\|_p$, $\|b\|_p$ (with $1 \leq p \leq \infty$) which explicitly depend on the initial data, see [7]. We point out that our derivation makes no explicit reference to the initial conditions (beyond the requirement $a(t, \mathbf{r}) \geq 0$, $b(t, \mathbf{r}) \geq 0$) and neither a scaling ansatz is needed. The bounds (1.6) reproduce the expected mean-field scaling $\bar{a}(t) \sim t^{-1}$. For the mathematically oriented reader we recall that in low dimensions $n < 2$, the description of the diffusive pair-annihilation processes through a more microscopic approach such as a master equation (where fluctuations are taken into account) leads to a different long-time behaviour $\bar{a}_{\text{micro}}(t) \sim t^{-n/2}$, which has also been observed experimentally for $n = 1$, see [8] for a recent review. This manifests once again the character of equations such as (1.1) as mean-field approximations.

The approach of $a(t, \mathbf{r})$ towards the mean density can be described as follows.

Corollary: Under the same conditions as in the theorem, there is a constant $K' \leq 1$ such that for times satisfying the conditions used in eq. (1.6)

$$\begin{aligned} \frac{1}{|\Omega|} \int_{\Omega} d\mathbf{r} \ (a(t, \mathbf{r}) - \bar{a}(t))^2 &\leq \frac{K'}{\lambda^2} \cdot t^{-2} \\ \frac{1}{|\Omega|} \int_{\Omega} d\mathbf{r} \ (b(t, \mathbf{r}) - \bar{b}(t))^2 &\leq \frac{K'}{2\mu} \cdot t^{-1} \end{aligned} \quad (1.7)$$

As an application, we consider the *pair-contact process* $2A \rightarrow \emptyset$, $2A \rightarrow 3A$ with single-particle diffusion (PCPD). We consider a domain $\Omega \subset \mathbb{R}^n$ with the boundary condition $\nabla a|_{\partial\Omega} = 0$. The mean-field reaction-diffusion equation is

$$\partial_t a(t, \mathbf{r}) = D \Delta a(t, \mathbf{r}) + \lambda a(t, \mathbf{r})^2 - \mu a(t, \mathbf{r})^3 \quad (1.8)$$

with constants $\lambda \in \mathbb{R}$ and $\mu > 0$. If the diffusion constant $D = 0$, $a = a(t)$ evolves for $\lambda > 0$ towards a steady-state density $a_\infty = \lambda/\mu$, while $a_\infty = 0$ for $\lambda \leq 0$. It is known that [4]

$$a(t) - a_\infty \sim \begin{cases} O(e^{-t/\tau}) & ; \text{ if } \lambda > 0 \\ t^{-1/2} & ; \text{ if } \lambda = 0 \\ t^{-1} & ; \text{ if } \lambda < 0 \end{cases} \quad (1.9)$$

as $t \rightarrow \infty$ and where $\tau > 0$ is a known constant. On the other hand, for a non-vanishing diffusion constant, we scale to $D = 1$ and have for the mean density

$$\lambda \bar{a}(t)^2 - \mu' \bar{a}(t)^3 \leq \partial_t \bar{a}(t) \leq \lambda' \bar{a}(t)^2 - \mu \bar{a}(t)^3 \quad (1.10)$$

for $\lambda \geq 0$ and

$$-|\lambda'| \bar{a}(t)^2 - \mu' \bar{a}(t)^3 \leq \partial_t \bar{a}(t) \leq -|\lambda| \bar{a}(t)^2 - \mu \bar{a}(t)^3 \quad (1.11)$$

for $\lambda \leq 0$, respectively and we can now let $|\Omega| \rightarrow \infty$, if we so desire. For $\lambda > 0$, there is an active steady-state with density $\lambda/\mu' \leq a_\infty \leq \lambda'/\mu$ but if $\lambda \leq 0$, one has $a_\infty = 0$. Furthermore, eq. (1.9) can be taken over. This proves the existence of a continuous steady-state transition at $\lambda_c = 0$ of the mean-field equation (1.8). Finally, the approach of $a(t, \mathbf{r})$ towards the mean density is according to

$$\frac{1}{|\Omega|} \int_{\Omega} d\mathbf{r} (a(t, \mathbf{r}) - \bar{a}(t))^2 \lesssim \begin{cases} O(e^{-2t/\tau'}) & ; \text{ if } \lambda > 0 \\ t^{-1} & ; \text{ if } \lambda = 0 \\ t^{-2} & ; \text{ if } \lambda < 0 \end{cases} \quad (1.12)$$

In $n = 1$ dimension, fluctuation effects create a very rich behaviour of the PCPD which is under active investigation, see [9] for a review.

In section 2, we recall some inequalities which are needed in the proofs. The upper bounds in (1.4,1.5) are derived in section 3 and in section 4, the lower bounds are obtained.

2 Mathematical background

We recall here some standard notations and some inequalities which will be needed in establishing the lower bound in (1.4,1.5). We shall work with the p -norms, for $1 \leq p < \infty$

$$\|u\|_p := \left(\int_{\Omega} d\mathbf{r} |u(\mathbf{r})|^p \right)^{1/p} \quad (2.1)$$

Denote by $L^p(\Omega)$ the space of (equivalence classes of) functions with $\|u\|_p$ finite. Here and in the following $\Omega \subset \mathbb{R}^n$. Furthermore, if Ω has a boundary, it is assumed to be sufficiently smooth throughout. The space $L^\infty(\Omega)$ is defined with respect to the supremum norm $\|u\|_\infty = \text{ess sup}_{\Omega} |u(\mathbf{r})|$. For derivatives, we use the multiindex notation $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}_0^n$ and where $|\alpha| := \alpha_1 + \dots + \alpha_n$. Then derivatives are denoted by

$$\partial^\alpha u = \frac{\partial^{|\alpha|} u}{(\partial r_1)^{\alpha_1} \dots (\partial r_n)^{\alpha_n}} \quad (2.2)$$

These derivatives can be taken to be weak (distributional) derivatives. The Sobolev space is

$$W^{k,p}(\Omega) := \{u \in L^p(\Omega), \partial^\alpha u \in L^p(\Omega) \text{ for all } |\alpha| \leq k\} \quad (2.3)$$

with its norm $\|u\|_{k,p} = \sum_{|\alpha| \leq k} \|\partial^\alpha u\|_p$. Finally, $C^1(\mathbb{R}_+; W^{k,p}(\Omega))$ is the space of functions which are continuously differentiable with respect to t for all times $0 \leq t < \infty$ and whose values at any given t are in $W^{k,p}(\Omega)$.

After these preparations, we can state some known results which we need later. The first one is the Gagliardo-Nirenberg inequality, see [6, 11].

Lemma 1. *For functions $u \in W^{k,p}(\Omega) \cap L^q(\Omega)$ with $1 \leq q \leq \infty$ and for any integer $0 \leq j < k$, there is a constant $C > 0$ such that*

$$\|\partial^j u\|_r \leq C \|u\|_q^{1-\theta} \|\partial^k u\|_p^\theta \quad (2.4)$$

where

$$\frac{1}{r} - \frac{j}{n} = \frac{1-\theta}{q} + \theta \left(\frac{1}{p} - \frac{k}{n} \right) \quad (2.5)$$

and if $1 \leq p < n/(k-j)$, then $j/k \leq \theta \leq 1$. On the other hand, if $1 \leq p = n/(k-j)$, only $j/k \leq \theta < 1$ is allowed.

We shall need two special cases of this. For $n \geq 2$, we set $p = r = 2$, $k = q = 1$. Then $j = 0$ and $\theta = n/(2+n) \in [\frac{1}{2}, 1)$. We get (here and in the following, we suppress the integration variable and write $\int_\Omega = \int_\Omega d\mathbf{r}$)

$$\int_\Omega |u|^2 \leq C_1 \left(\int_\Omega |u| \right)^{4/(2+n)} \left(\int_\Omega |\partial u|^2 \right)^{n/(2+n)} \quad (2.6)$$

Eq. (2.6) does not hold if $n < 2$. For $1 \leq n \leq 2$, we set $r = 2$, $p = q = k = 1$. Then $j = 0$, $\theta = n/2 \in [\frac{1}{2}, 1]$ and

$$\int_\Omega |u|^2 \leq C_2 \left(\int_\Omega |u| \right)^{2-n} \left(\int_\Omega |\partial u| \right)^n \quad (2.7)$$

where the constants $C_{1,2}$ equal C^2 from Lemma 1. It can be checked from dimensional analysis that C_1 and C_2 are independent of $|\Omega|$.

Next, we quote an inequality due to Nirenberg, see [6].

Lemma 2. *For $u \in W^{2,p}(\Omega)$, $p \geq 1$, there exists a constant $\varepsilon_0 = \varepsilon_0(p, \Omega)$ such that for any ε with $0 < \varepsilon < \varepsilon_0$, there is a positive constant $c = c(p, \Omega)$ such that*

$$\int_\Omega d\mathbf{r} |\partial u|^p \leq \frac{c}{\varepsilon} \int_\Omega d\mathbf{r} |u|^p + \varepsilon \sum_{|\alpha|=2} \int_\Omega d\mathbf{r} |\partial^\alpha u|^p \quad (2.8)$$

Finally, we quote Poincaré's inequality, see [11]. Let $B_R(0)$ be the ball of radius R around the origin.

Lemma 3. *For any $u \in W^{1,p}(B_R(0))$ with $1 < p < \infty$, there is a positive constant $C_p^{(p)}$ such that*

$$\int_{B_R(0)} d\mathbf{r} |u - \bar{u}|^p \leq C_p^{(p)} |B_R(0)|^{p/n} \int_{B_R(0)} d\mathbf{r} |\partial u|^p \quad (2.9)$$

and the mean value $\bar{u}$ is defined in analogy with (1.3).

It is sometimes desirable to consider spaces which are less restrictive than the spaces $L^p(\Omega)$. We define $\tilde{L}^p(\Omega)$ as the space of (equivalence classes of) functions such that $m_p(u) := |\Omega|^{-1} \|u\|_p^p$ is finite. For unbounded domains (e.g. $\Omega = \mathbb{R}^n$) a limit procedure must be used in the definition of $m_p(u)$. We also set

$$\widetilde{W}^{k,p}(\Omega) := \left\{ u \in \tilde{L}^p(\Omega), \partial^\alpha u \in \tilde{L}^p(\Omega) \text{ for all } |\alpha| \leq k \right\} \quad (2.10)$$

As an example, consider the function $f : \mathbb{R} \rightarrow \mathbb{R}$, $x \mapsto f(x) = f_0 \neq 0$. While $f \in \tilde{L}^p(\mathbb{R})$, since $m_p(f) = |f_0|^p$, clearly $f \notin L^p(\mathbb{R})$. Lemmas 2 and 3 readily extend to the space $\widetilde{W}^{k,p}(\Omega)$.

3 The upper bound

We briefly recall the proof of the upper bound in (1.4), following [1]. The mean density satisfies

$$\begin{aligned}\partial_t \bar{a} &= \frac{1}{|\Omega|} \int_{\Omega} d\mathbf{r} \Delta a - \frac{\lambda}{|\Omega|} \int_{\Omega} d\mathbf{r} a^2 \\ &= \frac{1}{|\Omega|} \int_{\partial\Omega} d\boldsymbol{\sigma} \cdot \nabla a - \frac{\lambda}{|\Omega|} \int_{\Omega} d\mathbf{r} a^2\end{aligned}\tag{3.1}$$

where $\boldsymbol{\sigma}$ is a normal vector to the boundary $\partial\Omega$. The first term describes the flux of particles through the boundary and vanishes either in the limit of large volumes $|\Omega| \rightarrow \infty$ or else if the boundary condition $\nabla a|_{\partial\Omega} = 0$ is imposed. Then $\partial_t \bar{a} = -\lambda \bar{a}^2 \leq -\lambda \bar{a}^2$ by the Cauchy-Schwarz inequality.

A similar argument works for triplet annihilation. By Hölder's inequality, $\partial_t \bar{b} = -\mu \bar{b}^3 \leq -\mu \bar{b}^3$.

4 The lower bound

In order to obtain the lower bound in (1.4), we recall from section 3 that $\partial_t \bar{a} = -\lambda \bar{a}^2$. The right-hand side is now estimated through the Gagliardo-Nirenberg inequality. We have to distinguish the cases $n \geq 2$ and $n \leq 2$ and obtain from eqs. (2.6) and (2.7)

$$\partial_t \bar{a} \geq \begin{cases} -\lambda C_1 |\Omega|^{-1} \left(\int_{\Omega} a \right)^{4/(2+n)} \left(\int_{\Omega} |\nabla a|^2 \right)^{n/(2+n)} & ; \text{ if } n \geq 2 \\ -\lambda C_2 |\Omega|^{-1} \left(\int_{\Omega} a \right)^{2-n} \left(\int_{\Omega} |\nabla a|^2 \right)^{n/2} & ; \text{ if } n \leq 2 \end{cases}\tag{4.1}$$

where for $n \leq 2$ the Cauchy-Schwarz inequality was used again. Next, we need an upper estimate for $\int_{\Omega} |\nabla a|^2$, which is provided by the following

Proposition: *For $u \in W^{2,p}(\Omega)$ there is a constant $c > 0$ and an ε_* such that $0 < \varepsilon_* < \infty$ and that for all $\varepsilon < \varepsilon_*$ one has*

$$\int_{\Omega} |\nabla u|^p \leq \frac{2c}{\varepsilon} \int_{\Omega} |u|^p\tag{4.2}$$

Proof: If $\sum_{|\alpha|=2} \int_{\Omega} |\partial^{\alpha} u|^p = 0$, the proposition holds true trivially, because of (2.8). We can thus suppose that $\sum_{|\alpha|=2} \int_{\Omega} |\partial^{\alpha} u|^p > 0$. Next, the function $f(x) := A/x + Bx$, where A, B are positive constants, has an absolute minimum at $x_* = \sqrt{A/B}$. For $x < x_*$, the first term dominates over the second and $f(x) < 2A/x$ for all $x < x_*$. We apply this to the inequality (2.8) of Lemma 2. The right-hand side is minimal if $\varepsilon = \varepsilon_*$, where

$$\varepsilon_* = \min \left(\left(\frac{c \int_{\Omega} |u|^p}{\sum_{|\alpha|=2} \int_{\Omega} |\partial^{\alpha} u|^p} \right)^{1/2}, \varepsilon_0 \right)$$

with the ε_0 of Lemma 2. Then the assertion follows. q.e.d.

The extension to $\widetilde{W}^{2,p}(\Omega)$ is immediate.

Therefore, setting $p = 2$ and appealing to dimensional analysis, there is a positive constant $K > 0$ such that

$$\begin{aligned}
\int_{\Omega} |\nabla a|^2 &\leq K|\Omega|^{-2/n} \int_{\Omega} a^2 \\
&= K|\Omega|^{-2/n} \int_{\Omega} [(a - \bar{a})^2 + 2\bar{a}(a - \bar{a}) + \bar{a}^2] \\
&= K|\Omega|^{1-2/n} \bar{a}^2 + K|\Omega|^{-2/n} \int_{\Omega} (a - \bar{a})^2 \\
&\leq 2K|\Omega|^{1-2/n} \bar{a}^2
\end{aligned} \tag{4.3}$$

From the eqs. (4.1) it follows

$$\partial_t \bar{a} \geq \begin{cases} -\lambda C_1 (2K)^{n/(2+n)} \bar{a}^2 & ; \text{ if } n \geq 2 \\ -\lambda C_2 (2K)^{n/2} \bar{a}^2 & ; \text{ if } n \leq 2 \end{cases} \tag{4.4}$$

This is exactly the form asserted in the theorem and we can identify the effective reaction rate

$$\lambda' := \begin{cases} \lambda C_1 (2K)^{n/(2+n)} & ; \text{ if } n \geq 2 \\ \lambda C_2 (2K)^{n/2} & ; \text{ if } n \leq 2 \end{cases} \tag{4.5}$$

Remark: the last bound in (4.3) might be improved by restricting Ω to a ball around the origin and applying the Poincaré inequality. If $C_p^{(2)} K < 1/2$, one gets $\int_{\Omega} |\nabla a|^2 < K/(1 - KC_p^{(2)}) |\Omega|^{1-2/n} \bar{a}^2$. For $n = 1$, the bounds might be further sharpened with the help of the inequality

$$\int_{\Omega} |\partial u|^p \leq c \varepsilon^{-\mu(p)} \left(\int_{\Omega} |u| \right)^p + \varepsilon \int_{\Omega} |\partial^2 u|^p \tag{4.6}$$

where $\mu(p) = -(p - 3 + 1/p)/p$ and which can be proven for $p \geq 1$ through a slight generalization of the proof [6] of the inequality (2.8) of Lemma 2. Since $\mu(p) > 0$ if $1 \leq p < p_c = (3 + \sqrt{5})/2$, the method of the proposition can for example be used in the $p = 1$ and $p = 2$ cases.

The lower bound in (1.5) is proved similarly. First, we set $k = q = 1$ and $r = 3$ in Lemma 1. Then $j = 0$. For $1 \leq n \leq 3/2$, we set $p = 1$, find $\theta = 2n/3 \in [2/3, 1]$ and

$$\int_{\Omega} |u|^3 \leq \Gamma_1 \left(\int_{\Omega} |u| \right)^{3-2n} \left(\int_{\Omega} |\partial u| \right)^{2n} \tag{4.7}$$

Next, for $3/2 \leq n \leq 3$, we set $p = 3/2$, find $\theta = 2n/(n+3) \in [2/3, 1]$ and

$$\int_{\Omega} |u|^3 \leq \Gamma_2 \left(\int_{\Omega} |u| \right)^{3(3-n)/(n+3)} \left(\int_{\Omega} |\partial u|^{3/2} \right)^{4n/(n+3)} \tag{4.8}$$

Finally, for $n \geq 3$, we set $p = 3$, find $\theta = 2n/(2n+3) \in [2/3, 1)$ and

$$\int_{\Omega} |u|^3 \leq \Gamma_3 \left(\int_{\Omega} |u| \right)^{9/(2n+3)} \left(\int_{\Omega} |\partial u|^3 \right)^{2n/(2n+3)} \tag{4.9}$$

Dimensional analysis shows that $\Gamma_{1,2,3}$ (which stand for C^3 of Lemma 1) are independent of $|\Omega|$. For the further analysis of (4.9), the proposition above states that there is a positive constant K_3 such that

$$\begin{aligned}
\int_{\Omega} |\nabla b|^3 &\leq K_3 |\Omega|^{-3/n} \int_{\Omega} b^3 \\
&= K_3 |\Omega|^{-3/n} \int_{\Omega} [\bar{b}^3 + 3\bar{b}(b - \bar{b})^2 + (b - \bar{b})^3] \\
&\leq K_3 |\Omega|^{-3/n} \left[|\Omega| \bar{b}^3 + 3\bar{b} \int_{\Omega} |b - \bar{b}|^2 + \int_{\Omega} |b - \bar{b}|^3 \right] \\
&\leq 5K_3 |\Omega|^{1-3/n} \bar{b}^3
\end{aligned} \tag{4.10}$$

Now, from $\partial_t \bar{b} = -\mu \bar{b}^3$, we obtain, using first eqns. (4.7,4.8,4.9), then the Cauchy-Schwarz inequality and the following consequence of Hölder's inequality $\int_{\Omega} |\nabla b|^{3/2} \leq |\Omega|^{1/4} (\int_{\Omega} |\nabla b|^2)^{3/4}$ and finally eqns. (4.3,4.10)

$$\begin{aligned}
\partial_t \bar{b} &\geq \begin{cases} -\mu \Gamma_1 |\Omega|^{-1} (\int_{\Omega} b)^{3-2n} (\int_{\Omega} |\nabla b|)^{2n} & ; \text{ if } 1 \leq n \leq 3/2 \\ -\mu \Gamma_2 |\Omega|^{-1} (\int_{\Omega} b)^{3(3-n)/(3+n)} (\int_{\Omega} |\nabla b|^{3/2})^{4n/(3+n)} & ; \text{ if } 3/2 \leq n \leq 3 \\ -\mu \Gamma_3 |\Omega|^{-1} (\int_{\Omega} b)^{9/(2n+3)} (\int_{\Omega} |\nabla b|^3)^{2n/(2n+3)} & ; \text{ if } 3 \leq n \end{cases} \\
&\geq \begin{cases} -\mu \Gamma_1 |\Omega|^{-1} (\int_{\Omega} b)^{3-2n} (\int_{\Omega} |\nabla b|^2)^n & ; \text{ if } 1 \leq n \leq 3/2 \\ -\mu \Gamma_2 |\Omega|^{-1+n/(n+3)} (\int_{\Omega} b)^{3(3-n)/(3+n)} (\int_{\Omega} |\nabla b|^2)^{3n/(3+n)} & ; \text{ if } 3/2 \leq n \leq 3 \\ -\mu \Gamma_3 |\Omega|^{-1} (\int_{\Omega} b)^{9/(2n+3)} (\int_{\Omega} |\nabla b|^3)^{2n/(2n+3)} & ; \text{ if } 3 \leq n \end{cases} \\
&\geq \begin{cases} -\mu \Gamma_1 (2K)^n \bar{b}^3 & ; \text{ if } 1 \leq n \leq 3/2 \\ -\mu \Gamma_2 (2K)^{3n/(n+3)} \bar{b}^3 & ; \text{ if } 3/2 \leq n \leq 3 \\ -\mu \Gamma_3 (5K_3)^{2n/(2n+3)} \bar{b}^3 & ; \text{ if } 3 \leq n \end{cases}
\end{aligned} \tag{4.11}$$

which is the form asserted in (1.5). We identify the effective reaction rate

$$\mu' := \begin{cases} \mu \Gamma_1 (2K)^n & ; \text{ if } 1 \leq n \leq 3/2 \\ \mu \Gamma_2 (2K)^{3n/(n+3)} & ; \text{ if } 3/2 \leq n \leq 3 \\ \mu \Gamma_3 (5K_3)^{2n/(2n+3)} & ; \text{ if } 3 \leq n \end{cases} \tag{4.12}$$

This completes the proof of the theorem. q.e.d.

The bounds (1.6) are established by direct integration, since $\partial_t \bar{a}(t) \geq -\lambda \bar{a}(t)^2$ implies $\bar{a}(t) \geq a_0/(1 + a_0 \lambda t)$ from which the assertion is immediate, under the stated condition on t and for $a_0 > 0$. The bounds for $\bar{b}(t)$ follow similarly.

We now prove the corollary. First, $|\Omega|^{-1} \int_{\Omega} (a - \bar{a})^2 \leq |\Omega|^{-1} \int_{\Omega} \bar{a}^2 = \bar{a}^2 \leq (\lambda t)^{-2}$. Next, we try to improve this bound by letting $\Omega = B_R(0)$ be a ball around the origin and apply Poincaré's inequality

$$\frac{1}{|\Omega|} \int_{\Omega} (a - \bar{a})^2 \leq C_P^{(2)} |\Omega|^{-1+2/n} \int_{\Omega} |\nabla a|^2 \leq 2K C_P^{(2)} \bar{a}^2$$

and therefore $K' = \min(1, 2K C_P^{(2)})$. The estimates for b are obtained similarly. q.e.d.

Remark: for the PCPD with $D = 1$, eqns. (1.10,1.11) are obtained by applying the results of the theorem separately to the two terms on the right-hand side of eqn. (1.8). Eq. (1.12) is obtained by replacing $a \mapsto a - a_{\infty}$ in the corollary and then using (1.9). Let $a_t(\mathbf{r}) := a(t, \mathbf{r})$. For sufficiently long times, $a_t \in \widetilde{W}^{2,3}(\mathbb{R}^n)$, but if $\lambda > 0$, $a_t \notin W^{2,3}(\mathbb{R}^n)$.

Acknowledgments

I thank R. Cherniha, H. Spohn, M. Struwe and P. Wittwer for useful conversations or correspondence and the organizers of the XIVth International Conference on Mathematical Physics in Lisbon, especially J.-C. Zambrini, for providing the stimulating atmosphere where the ideas presented here formed. The warm hospitality of the Centro de Física Teórica e Computacional (CFTC) of the Universidade de Lisboa is gratefully acknowledged.

References

- [1] D. ben-Avraham, M.A. Burschka and C.R. Doering, J. Stat. Phys. **60**, 695 (1990).
- [2] G. van Baalen, A. Schenkel and P. Wittwer, Comm. Math. Phys. **210**, 145 (2000).
- [3] J. Bricmont, A. Kupiainen and G. Lin, Comm. Pure Appl. Math. **47**, 893 (1994).
- [4] E. Carlon, M. Henkel and U. Schollwöck, Phys. Rev. **E63**, 036101 (2001).
- [5] S. Cornell and M. Droz, Phys. Rev. Lett. **70**, 3824 (1993).
- [6] A. Friedman, *Partial differential equations*, Holt, Rinehart and Winston (New York 1969); sects. 8, 10.
- [7] A. Gmira and L. Veron, J. Differential Equations **53**, 258 (1984).
- [8] M. Henkel, in A. Kundu (ed), *Classical and Quantum Nonlinear Integrable Systems: Theory and Application*, (IOP, Bristol 2003); ch. 10 (cond-mat/0303512).
- [9] M. Henkel and H. Hinrichsen, ‘‘The non-equilibrium phase transition of the pair-contact process with diffusion’’, in preparation.
- [10] A. Schenkel, P. Wittwer and J. Stubbe, Physica **D69**, 135 (1993).
- [11] J. Shatah and M. Struwe, *Geometric wave equations*, Courant Lecture Notes in Mathematics Vol. 2, American Mathematical Society (New York 2000); ch. 3.
- [12] J. Smoller, *Shock waves and reaction-diffusion equations*, Springer (Heidelberg 1983).
- [13] S.V. Zelik, Comm. Pure Appl. Math. **56**, 584 (2003).